Index

page numbers in *italics* denote figures and exhibits; page numbers followed by t denote tables

A

Abscess, 109t
Accreditation, 2, 50, 51t, 224–225, 227
Acid-fast stain, 9
Acinetobacter, 10t
ACIP. *See* Advisory Committee on Immunization Practices (ACIP)
Acquired immunodeficiency syndrome (AIDS). *See* Human immunodeficiency virus (HIV)
Actinomycosis, 109t
Adenovirus, 10t, 109t, 112t
Advisory Committee on Immunization Practices (ACIP), 197–198
Agency for Health Care Policy and Research (AHRQ)
  grants related to home care safety, 221
  guidelines for treatment of pressure ulcers, 23–24
AIDS. *See* Human immunodeficiency virus (HIV)
Airborne Isolation, 101–103
  diseases transmitted by airborne droplet nuclei, 101, 102–103
  patient placement for, 101
  during patient transpost, 103
  personal protective equipment for, 102
  prevention of exposure to, 102
Airborne transmission of disease, 12
  respiratory protection from, 82, *83–84*
  for tuberculosis, 82
Alcohols, as antiseptic agents, 18t, 123
  fire hazards and storage of alcohol-based hand rubs, 21–22
  instructions for use of, 21t
Allergic reactions to latex, 80–81, 81t
Amebiasis, 109t
American Academy of Pediatrics, on vaccines for children, 198
American Association of Respiratory Care, 30
American Hospital Association (AHA), surveillance of nosocomial infections, 2
American Society of Health-System Pharmacists (ASHP), 50
Animal-assisted activities (AAA)/animal-assisted therapy (AAT), 68–69
  animals that should not be used for, 69
  differences between, 69
  guidelines when a program is established, 70–71
  hand hygiene and cleaning procedures, 71
  transmission of zoonotic diseases and, 69–71, 70t
Animals
  diseases carried by, 70t
  transmission of infection by, 9, 22
Anthrax, 109t
Antibiotic-associated colitis, 109t
Antigenic variation, 8t, 9
Antimicrobial agents, 9, 17–18, 18t
  resistance to (*See* Multidrug resistant organisms (MDROs); *specific organisms*)
Antiseptic, 122–123
APIC. *See* Association for Professionals in Infection Control (APIC)
Arthropod-borne viral diseases, 109t, 110t
Ascariasis, 110t
Aseptic technique
  defined, 23
  preparation of medication in the home, 52
Aspergillosis, 110t
Aspergillus niger, 11t, 12
Aspiration, preventing respiratory infection caused by, 33
Assessment
  health assessment and immunizations, 197
  initial assessment and implementation of isolation precautions, 105–106
  of patient and home environment, 22

Assessment *(continued)*
population assessment, 168–169, *169*
of tuberculosis, 202–205
Association for Professionals in Infection Control (APIC)
APIC Curriculum for Infection Control, 2
domains for infection control practice, 2
guideline for Sterilization and Disinfection, 121
recommendations for infection control programs, 222
Avian influenza, 110t

B

Babesiosis, 110t
Bacillus anthracis, 72t
Bacillus cereus, 72t
Bacteria, 10t
bacilli, 9
cocci, 9
Gram negative, 9, 90
Gram positive, 9, 90
resistance in DNA of, 91
Bacterial diseases
associated with zoonotic transmission, 70t
foodborne, 72–75t
Biohazard symbol, *140*
Bioterrorism, agents of, 208–209
Blastomycosis, 110t
Bleach solution, 123, 124, 126
for linens and laundry, 128
preparing, 125, *125*
storing, 125
Blepharitis, 119t
Bloodborne pathogens
defining and managing exposures to, 209–212
OSHA regulations, 79, 123, 124, 131, 200–201, *201*
Bloodborne transmission of disease, 7
Blood cultures, for bacteria, 90
Blood pressure cuffs, cleaning and disinfection of, 125
Blood spills in the home, cleaning, 140
Blood storage for home transfusions
in the patient's home, 50
during transport, 49–50
Bloodstream infection (BSI), 37–40
definitions of for surveillance purposes, 172, 172t
incidence density, central line-associated bloodstream infection, 181, *182*
line listing for IV catheter-related bloodstream infections, 179, *180*
organisms causing, 181, *181*
performance indicators from CDC guidelines for IV catheter-related bloodstream infections, 179, *180*
summary of, 181, 181t
Botulism, 110t, 112t
Breast milk. *See also* Infants
storage of, 68
thawing of frozen milk, 68
wearing of gloves and contact with, 80
Breathing circuits, 30
Bronchiolitis, 110t
Broth dilution, 90
Broviac catheter, 40, 48
Brucella abortus, 72t
Brucella melitensis, 72t
Brucella suis, 72t
Brucellosis, 110t
BSI. *See* Bloodstream infection (BSI)
Burkholderia cepacia, 90

C

Campylobacter gastroenteritis, 110t
Campylobacter jejuni, 10t, 72t
Campylobacter spp., 112t
Candida albicans, 11t, 27
Candidiasis, 110t
Cardiopulmonary resuscitation (CPR), use of resuscitation mask, 85
Carpet surfaces, decontamination of, 140
Carriers, for transmission of disease, 7, 192
Catheter-associated urinary tract infection (CAUTI), 26–27. *See also* Urinary tract care; Urinary tract infection (UTI)
surveillance data, 180–181, 180t, 182, 183
Catheter site care
injection cap changes, 46
midline and central venous catheter dressing changes, 45–46
skin preparation, 45
Cat-scratch fever, 110t
Causative agents of infection, 8–9, 8t, 10–11t
CAUTI. *See* Catheter-associated urinary tract infection (CAUTI)
Cellulitis, 110t
Center for Disease Control (CDC)
Epidemiologic Intelligence Service (EIS), 192–193
guidelines
for changing of breathing circuits, 30
Draft Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings, 51t, 57–59t, 62–63, 89, 93–96, 95t
Guideline for Hand Hygiene in Health-Care Settings, 16, 16t
Guideline for the Prevention of Catheter-Associated Urinary Tract Infections, 27
"Guidelines for Preventing the Transmission of *Mycobacterium tuberculosis* in Health-Care Facilities," 202
for HIV postexposure prophylaxis, 219–220

performance indicators from selected guidelines, *226*, 227
for prevention of health care acquired infections, 3–4, 3t
for respiratory protection, 82
on vaccines for children, 198
study on bloodborne exposures among home care staff members, 209–210
training for infection control, 2
Central venous catheters
for children, 43, 48
dressing changes for, 45–46
infections related to, CDC guidelines for prevention of, 63
nontunneled, 39–40
peripherally inserted, 40 (*See also* Peripherally inserted central catheter (PICC))
tunneled, 40
Chain of infection, 8, 8f
Chancroid, 110t
Chemical sterilant, 123
Chickenpox. *See* Varicella
Childhood diseases
Airborne Precautions for, 104–105
vaccine-preventable, 197–208, *198*
Children. *See* Pediatric patients
Chlamydia trachomatis, 110t
Chlorhexidine, as antiseptic agent, 18t
Cholera, 110t, 112t
epidemic, London (1849), 163
Cidex, 126
Cleaning, meaning of the term, 121–122
Cleaning and disinfection, 121–130
bleach solution, 123, 124, 126
preparing, 125, *125*
storing, 125
categorization of home care equipment, 121, 122t
catheters, 127t
critical items, 121, 122t, 126, 128
definition of terms
antiseptic, 122–123
chemical sterilant, 123
cleaning, 121–122
decontamination, 122
disinfectant, 123
germicide, 122–123
sterilization, 123
dishes, glasses, cups, and eating utensils, 129
effectiveness of disinfection process, determining, 124
enteral feeding equipment and supplies, 34
environmental surfaces, 128
EPA registered products for, 123–124
hinged instruments, 127t
identification and usage of medical devices in the home, 121
inner tracheal cannula, 31–32
intermittent urethral catheters, 29
lensed instruments, 127t
levels of disinfection, 123
high-level, 123
intermediate-level, 123–124
low-level, 123
linens and laundry, 128
of medical equipment before reuse, 26, 126
noncritical items, 121, 122t
commonly used items, 125
laboratory supplies, 125–126
methods of sterilization and disinfection, 127t
provided to patients, 126
for pediatric patients
cleaning of surfaces likely to be touched by, 65
clothing and linen, 66
diaper changing area, 66
"potty training" equipment, 66
resource guidelines for, 121
respiratory equipment and supplies, 31–32, 32
rubber tubing, 127t
semicritical items, 121, 122t, 126
Spaulding's scheme, 121
storage of equipment and supplies
in the home care or hospice facility, 129
in patient's home, 130
during patient transport, 129–130
surfaces, 127t
thermometers, 127t
tracheal cannula, 31–32
tracheal suction catheters, 31
urine collection and tubing bags, 29
use of personal protective equipment, 124
Clean technique, and sterile technique, comparison of, 23
Clinical Practice Guidelines, treatment of pressure ulcers, 23–24
Clinical service groups (CSGs), 225
Clostridium botulinum, 72t, 110t
Clostridium difficile, 10t, 110t, 111t, 112t
Clostridium perfringens toxin, 72t, 111t, 112t
Clostridium tetany, 10t
Clostridium welchii, 112t
Clothing
cleaning and disinfection for, 66
Contact Precautions for, 105
protective (*See* Personal protective equipment (PPE))
CMV. *See* Cytomegalovirus (CMV)
Coccidioidomycosis, 111t
Colitis, antibiotic-associated, 109t
Colonization, 8, 173

Colorado tick fever, 111t
Communication, during an outbreak situation, 193
Community-acquired infections, 188
Comprehensive Accreditation Manual for Home Care (JCAHO), 50, 51t
Condom catheter drainage, 27
Conjunctivitis, 111t, 119t
  as indication for work restrictions, 212t, 216
Contact, transmission of disease by, 11–12
Contact Precautions, 95–96
  discontinuing, guidelines for, 105
  diseases or conditions that require, 104–105
  glowns, gloves, and handwashing for, 105
  during patient transport, 105
Cornybacterium diphtheriae, 10t
Corona virus associated with SARS (SARS-CoV), 111t
Costs, of health care-associated infection, 4
Coxsackie virus, 10t, 111t
CPR. *See* Cardiopulmonary resuscitation (CPR)
Creutzfeldt-Jacob disease, 111t
Croup, 111t
Cryptococcosis, 111t
Cryptosporidiosis, 111t
Cryptosporidium spp., 112t
Crytosporidium parvum, 78
CSGs (clinical service groups), 225
Cysticerosis, 111t
Cytomegalovirus (CMV), 10t, 111t
  concerns for pregnant staff members regarding, 216, 217t
  as indication for work restrictions, 212t

D

Data collection methods. *See also* Infection surveillance
  defining denominators, 174, 176t
  description of the infection, 173
  design of the form, 174
  determining what to record, 173
  identifying home care-acquired infections, 177
    confirmed home care-acquired infection report, *178–179*
    infection screening report, *175–176*
Decontamination, 122
Decubitus ulcer, 111t
Dengue fever, 111t
Department of Health and Human Services (HHS), 4
Department of Transportation (DOT), requirements for medical waste transport, 139
Dermatomycosis, 118t
Dermatophytosis, 118t
Dialysis, transfer of MDROs and, 92
Diaper changing, *65*, 65–66
Diarrhea, 111t
  as indication for empiric precautions, 106t
  as indication for work restrictions, 212t, 215–216, 216
Diphtheria, 111t
  as indication for work restrictions, 212t
Diphtheria vaccine, 200
Disinfectant, 123
Disk diffusion, 90
Donovanosis granuloma venereum, 113t
Draft Isolation Guideline (CDC), 100
Dressings
  catheter dressing changes, 45–46
  changing, procedures for, 24–25
Drop box, for disposal of medical sharps, 156
Droplet Precautions, 103
  diseases or conditions that require, 103
  indications for use of, 103
  patient placement for, 104
  during patient transport, 104
  personal protective equipment for, 103–104
Droplet spread/droplet transmission, 11

E

Ebola, 103, 111t
Echinococcosis, 111t
Echovirus, 10t, 111t
Eczema vaccinatum, 119t
Education
  home care and hospice staff, 2, 85, 86t, 96, 223–224, *224*
  on location and use of personal protective equipment, 85, 86t
  patient and caregiver, home infusion therapy, 53–54
  patient and family
    isolation precautions, 107
    prevention of respiratory tract infection, 33
    prevention of urinary tract infection, 29–30
    related to enteral therapy, 34
    related to wound care, 26
EHEC (enterohemorrhagic *escherichia coli*), 72t
EIS (Epidemiologic Intelligence Service), 192–193
Elderly persons, risk for infection, 13
Encephalitis, 111t, 113t
Encephalomyelitis, 111t
Endocarditis, 37
Endometritis, 111t
  definitions of for surveillance purposes, 172t
Engineering controls (OSHA), 131
Enteral feeding equipment and supplies, cleaning and disinfection of, 34
Enteral therapy, 34
Enterobacter
  as cause of BSIs, 38

as cause of urinary tract infections, 27
Enterobacter aerogenes, 10t
Enterobiasis, 111t
Enterococci, 27, 111t
Enterocolitis, 111t, 118t
Enterohemorrhagic *escherichia coli* (EHEC), 72t
Enterotoxigenic *escherichia coli* (ETEC), 72t
Enteroviral infections, 112t
as indication for work restrictions, 212t
Environmental control, Standard Precautions for, 101t
Environmental Protection Agency (EPA)
medical sharps disposal, 156
recommendations for medical waste management plan, 131
registered products for cleaning and disinfection, 123–124
Environmental surfaces
cleaning and disinfection of, 128
touched by children, cleaning and disinfection of, 65
Epidemiologic Intelligence Service (EIS), 192–193
Epidemiologic triangle, 7, *8*
Epidemiologist, 2
Epidemiology, of nosocomial infection, 2–3
Epidural catheter, port, or pump, 49
Epiglottitis, 112t
Epstein-Barr virus infection, 112t
Equipment and devices. *See* Medical devices
Erythema, 112t, 120t
Escherichia coli, 7, 10t, 16
0157:H7, 71, 72t, 112t
as cause of urinary tract infections, 27
enterohemorrhagic (EHEC), 72t
enterotoxigenic (ETEC), 72t
gastroenteritis, 112t
shigatoxin-producing *E. coli* (STEC), 72t
ETEC (enterotoxigenic *escherichia coli*), 72t
Exanthems, 106t
Exit-site infections, 37
definitions of for surveillance purposes, 172–173
Eye protection, 82, *83–84*

F

Face shields, 82, *83–84*
Feeding pediatric patients
preparing infant formula
method, 67
storage of formula, 67–68
storage of human milk, 68
terminal sterilization, 67
thawing frozen milk, 68
preparing infants' bottles, 66–67
Females, risk for urinary infections, 13
Fetal vaccinia, 119t
Filters, in-line, home infusion therapy, 46–47
Fire hazards, of alcohol-based hand rubs, 21–22
Fleming, Alexander, 89
Food and Drug Administration (FDA), rental of reusable equipment, 126
Foodborne illness, 71, 72–75t. *See also* Meals for patients
Food poisoning, 112t
Foods, associated with foodborne illnesses, 73t, 75t
Formula. *See* Infants, feeding
Fungal diseases, associated with zoonotic transmission, 70t
Fungal infections, 11t, 118t
Furunculosis, 112t

G

Gangrene, 112t
Gastroenteritis, 110t, 112t
Gender, risk for infection and, 13
German measles, 107, 117t
Germicide, 122–123
Giardia lamblia, 112t
Giardiasis, 113t
Gloves, protective
for Contact Precautions, 105
latex allergies and, 80–81
prevention of in the workplace, 81
products containing latex, 81t
microbial contamination of, 79
Standard Precautions for, 101t
types of, 80
when to change or remove, 80
when to wear, 80
Goggles, 82, *83–84*
Gonococcal ophthalmia neonatorum, 113t
Gonorrhea, 113t
Gowns, protective, 82, *83–84*
for Contact Precautions, 105
Standard Precautions for, 101t
use of with patients with MDROs, 96
Gram's stain, 9, 90
Granuloma inguinale, 113t
Guideline for Environmentl Infection Control in HealthCare Facilities, 121
Guideline for Hand Hygiene in Health-Care Settings, 16, 16t
Guideline for the Prevention of Catheter-Associated Urinary Tract Infections, 27
Guideline for the Prevention of Catheter-Associated Urinary Tract Infections, 3
"Guidelines for Preventing the Transmission of *Mycobacterium tuberculosis* in Health-Care Facilities," 202
Guillan-Barré syndrome, 113t

H

Hand, foot, and mouth disease, 113t
Hand hygiene, 2, 14, 15–22
  before and after catheter maintenance, 28
  after removing protective gloves, 79
  antiseptic hand rubs, 20–21, 21t
  bar soap *vs.* liquid soap, 18
  behavioral aspects of, 19–20
  for Contact Precautions, 105
  effect of resident and transient microorganisms, 15–16
  facilities for, 18–19
  fire hazards and storage of alcohol-based hand rubs, 21–22
  hand creams, for contact dermatitis, 19
  indications for handwashing and hand antisepsis, 17, *17*
  indicators adapted for home and hospice care, *226*
  instructions for washing hands with soap and water, 20
  length of fingernails, considerations for staff, 19
  performance indicators from CDC guidelines, *226*
  plain soaps *vs.* antiseptic agents, 17–18, 18t
  purchasing hand hygiene products, 19
  side effects of antimicrobial soaps, 19
  Standard Precautions and, 101t
  supplies needed by home care staff, 20
  terminology definitions, 16, 16t
  using patient's hand hygiene supplies, 20
  waterless antiseptic agents, 17–18, 19
Hantavirus pulmonary syndrome (HPS), 22, 113t
HAV. *See* Hepatitis A virus (HAV)
Hawthorne effect, 192
HBV. *See* Hepatitis B virus (HBV)
HCV. *See* Hepatitis C virus (HCV)
Health Care Financing Administration (HCFA), reimbursement for catheter insertion and replacement, 27
Healthcare Infection Control Policy Advisory Committee (HICPAC), 3–4, *226*, 227
Helicobacter pylori, 113t
Hemodialysis catheters, 40–41
Hemophilus influenzae type b disease, Droplet Precautions for, 103, 112t
Hepatitis A virus (HAV), 10t, 113t
  concerns for pregnant staff members regarding, 217t
  defining and managing exposures to, 207
  as indication for work restrictions, 212t
Hepatitis B immune globulin (HBIG), 210
Hepatitis B surface antigen (HBsAG), 210
Hepatitis B vaccine, 195, 200–202, *201*
  testing for anti-HBs after vaccination, 201–202
Hepatitis B virus (HBV), 7, 10t, 113t, 209, 210
  concerns for pregnant staff members regarding, 217t
  defining and managing exposure to, 209
  as indication for work restrictions, 212–213t
Hepatitis C virus (HCV), 7, 10t, 113t
  concerns for pregnant staff members regarding, 217t
  defining and managing exposure to, 209, 210
  as indication for work restrictions, 213t
Hepatitis D virus, 113t
Hepatitis E virus, 113t
Hepatitis G virus, 113t
Herpangina, 113t
Herpes simplex virus, 10t, 113–114t
  concerns for pregnant staff members regarding, 217t
  as indication for work restrictions, 213t
  type 2, 10t
Herpes simplex virus (HSV), as indication for work restrictions, 215–216
Herpes zoster, 114t
  concerns for pregnant staff members regarding, 217t
  as indication for work restrictions, 214t
HHS (Department of Health and Human Services), 4
Hickman catheter, 40
HICPAC) (Healthcare Infection Control Policy Advisory Committee), 3–4
Hinged instruments, cleaning and disinfection of, 127t
Histoplasmosis, 114t
HIV/AIDS. *See* Human immunodeficiency virus (HIV)
Home care-acquired infection
  definitions for, 165
  developing, 171–173, 172t
  measuring the incidence, 4
  outbreak investigations, 188 (*See also* Outbreak investigations)
  results of home care-acquired infection rates, 165–166, 165t
  sites of, 170, *170*
  surveillance (*See* Infection surveillance)
Home environment, assessment of, 22
Home infusion therapy, 37–63
  administration of total parenteral nutrition (TPN), *47*, 48
  blood storage for home transfusions
    in the patient's home, 50
    during transport, 49–50
  catheter site care
    injection cap changes, 46
    midline and central venous catheter dressing changes, 45–46
    skin preparation, 45
  causes of IV catheter-related infections, 38, *39*
  culturing for suspected infection related to, 47–48
  distant and local infections related to, 37
  etiology of healthcare-associated BSIs, 38
  flushing the catheter, 47
  incidence of infection with, 37–38
  indicators adapted for home and hospice care, *226*
  infection control in pharmaceutical services, 50–53

medication storage, 52
multidose vials, 53
nursing care and administration of parenteral medications, 53
pharmacy sterile compounding requirements (USP and ASHP), 50, 51t, 57–59
preparation of parenteral medication in the home, 52
storage and transport of parenteral medications, 52
USP-NF Chapter 797 requirements, 50, 51t, 57–59
inserting catheter for, 43
needleless intravascular devices, 44
nonvascular access devices, 48–49
epidural catheter, port, or pump, 49
intrathecal catheter, port, or pump, 49
ventricular, 48, 49
overview of, 37–38
patient and caregiver education, 53–54
pediatric patients, 48
preventing catheter-related infections, 41
removing IV access devices, 44
replacing administration sets, 44
replacing IV access devices, 43–44
replacing IV solutions, 44–45
risk factors associated with catheter-related infections during TPN, *47*
selecting a catheter insertion site, 42–43
midline venous access and PICC site selection, 42
peripheral venous access site, 41–42
surveillance of infections related to, 171–172
thrombogenic catheter material, 43
types of catheters and devices
central venous catheters
nontunneled, 39–40
peripherally inserted central catheter (PICC), 40
tunneled, 40
hemodialysis catheters, 40–41
implanted ports, 41
midline catheter, 39
peripheral venous catheters, 38–39, *39*
use of in-line filters, 46–47
USP-NF 797 chapter, summary of, 57–59t
Host specificity, 8t, 9
Host susceptibility to infection, 12–14
extrinsic risk factors, 13–14
intrinsic risk factors, 12–13
HPS (hantavirus pulmonary syndrome), 22
Human immunodeficiency virus (HIV), 10t, 109t, 114t
CDC recommendations for postexposure prophylaxis, 219–220
concerns for pregnant staff members regarding, 217t
defining and managing exposure to, 210
incidence of infection for patients on home infusion therapy, 37–38
as indication for work restrictions, 213t
from occupational exposure, 195
Human reservoirs of infection, 9
Hymenolepsis nana, 118t

I

ICN (infection control nurse), 2
ICP. *See* Infection control practitioner (ICP)
Immune status, risk for infection and, 13
Immunizations. *See also specific diseases*
for HBV, 195
for home care staff, 197
paying for testing and immunization, 198
preemployment health survey/immunization status, 198, *199*
for influenza, 22, 33–34
for pneumonia, 22
recommendations for, 197
vaccine-preventable diseases, 197–202, *198*
Impetigo, 114t
Implanted ports, 41
infections related to, CDC guidelines for prevention of, 63
Incubation period for infection, 173
Infants
diapering, *65,* 65–66
cleaning and disinfecting diaper changing area, 66
feeding
preparing infant formula
method, 67
storage of formula, 67–68
storage of human milk, 68
terminal sterilization, 67
thawing frozen milk, 68
preparing infants' bottles, 66–67
risk for infection, 13
Infection
acute signs and symptoms of, 7
chain of, 8, 8f
chronic, 7
closed-cavity, 110t
device-related, 9, 13–14
extrinsic risk factors, 13–14
intrinsic risk factors, 12–13
normal flora, 7–8
postpartum, 1
reducing the risk for, 12
sites of home care-acquired infection, 170, *170*
subclinical, 7
types of, 7
Infection control nurse (ICN), 2
Infection control practitioner (ICP), 164, 188–193, 224, *224*
occupational health services by, 196

Infection control programs, 221–227
developing and maintaining
agencies focusing on patient safety in home care, 221
approaches to, 221
clinical service groups, 225
components of occupational/infection control policies, *223*
content for an orientation program, *223*
defining the program, 221–224
designation of an infection control practitioner (ICP), 224, *224*
functions of, *222*
importance of programs, 221
indicators adapted for home and hospice care, *226*
JCAHO accreditation and infection control, 224–225, 227
JCAHO safety goals for home care, 221, *222*
maintaining and advancing the program, *226,* 227
patient care practices, policies, and procedures, *222*
performance indicators from CDC guidelines, *226*
priority focus areas, 225
staff education, 223–224, *224*
surveillance program, 223
development of in the United States, 1–2
essential domains for, 2
history of, 1
hospital accreditation standards for, 2
status of, in home care and hospice, 3–4, 3t
Infection control supplies, 87, *87*
Infection surveillance, 2, 163–184
aggregation and analysis of infection data, 177, 179–183
central line-associated UTI data by quarter, 180–181, 180t, 182
formulas for calculation of infection rates, 179, *180*
line listing for IV catheter-related bloodstream infections, 179, *180*
organisms causing bloodstream infections, 181, *181*
proportions summary of home care-acquired infection by site, 181, *181*
summary of bloodstream infections, 181, 181t
cultures to identify patients with MDROs, 92–93
cumulative incidence post operative pneumonia, 181, *182*
CVC-associated infections, 166–167, 167t, 168
data collection methods
defining denominators, 174, 176t
description of the infection, 173
design of the form, 174
determining what to record, 173
identifying home care-acquired infections, 177
confirmed home care-acquired infection report, *178–179*
infection screening report, *175–176*
period of surveillance, 174
retrospective *vs.* concurrent data collection, 174, 177
risk factors by site, 173, 174t
defined, 163
definitions for home care-acquired infections, 165
developing, 171–173, 172t
endemic infection, 164
history of, 163–164
identifying potential causes of infection, 164
incidence density, central line-associated bloodstream infection, 181, *182*
incidence of healthcare-associated infections, study of, 165
measuring the incidence of home care-acquired infections, 4
nosocomial infection and, 163–164
ongoing measurement, 183
population assessment, 168–169, *169*
principles and methodology of, 163
reasons for, 164–168
results of home care-acquired infection rates, 165–166, 165t
selection of outcomes or processes for measurement, 169–171, *170*
software for surveillance, 184
steps in developing a surveillance program, 168, *168,* 184
symptomatic UTI infection, 166, 166t, 168
use of data for improvement of patient care, 183
validating surveillance data, 183–184
Infectious disease process
agent, host, and environment, 7
causative agent, 8–9, 8t
chain of infection, 8, 8f
common pathogenic agents, 10–11t
epidemiologic triangle, 7, 8f
extrinsic risk factors, 13–14
impact of home care and hospice staff, 14
intrinsic risk factors, 12–13
mode of transmission, 11–12
normal flora, 7–8
portal of entry, 11
portal of exit, 11
reservoir of infection, 9
risk of infection, reducing, 12
susceptible host, 12–14
Infectious mononucleosis, 114t
Infective dose, 8, 8t
Inflammation, 173
Influenza, 100, 114t
A2 and B, 9, 11t
prevention and control of, 33–34
Influenza vaccine, 22, 205
Infusion Nurses Society's Standards of Practice, 43–44

Injection cap changes, 46
In-line filters, home infusion therapy, 46–47
Inner tracheal cannula, cleaning and disinfection of, 31–32
Institute of Medicine (IOM), report on health care delivery, 221, 224
Intensive care units (ICUs), risk of MDROs in, 91–92
Intermittent urethral catheters, cleaning and disinfection of, 29
International Health Care Worker Safety Center, 134
Interrater reliability, 171
Intrathecal catheter, port, or pump, 49
Invasiveness, 8–9, 8t
In vitro testing, 90
In vivo testing, 90
Iodophors, 18t, 19, 123
IOM (Institute of Medicine), report on health care delivery, 221
Iritis, 119t
Irrigation solutions
  maintenance of in the home, 25, 25t
  preparation of, 25–26
Isolation Precautions, 99–120
  CDC guidelines for, 51t, 57–59t, 62–63, 89, 91–94, 93–96, 93t, 95t, 100
  chain of infection and, 99–100
  Contact Precautions, 95–96
  development of, 2
  goals of in home care, 99
  indications for empiric precautions to prevent transmission, 106t
  initial assessment and implementation of, 105–106
  patient and family education for, 107
  patient confidentiality and, 106–107
  performance indicators from CDC guidelines, *226*
  principles, application of, 99, 100
  Standard Precautions, 100
  storing wound care supplies, 25
  transmission-based precautions, 100–105
    Airborne Isolation, 101–103
    Contact Precautions, 104–105
    droplet precautions, 103–104
    Protective Environment (PE), 100–101
    types of and indications for, 100–101
  type and duration of, for selected infections and conditions, 109–120t
Isolation Techniques for Use in Hospitals, 2

J

Joint Commission on Accreditation of Healthcare Organizations (JCAHO), 2
  accreditation and infection control, 224–225, 227
  home care standards, 50, 51t
  safety goals for home care, 221, *222*

K

Kawasaki syndrome, 114t
Keratitis, 119t
Keratoconjunctivitis (EKC), as indication for work restrictions, 216
Klebsiella
  as cause of urinary tract infections, 27
  in ICU patients, 91
Klebsiella oxytoca, 92
Klebsiella pneumoniae, 10t
Koch, Robert, 1

L

Laboratory supplies, cleaning and disinfection of, 125–126
Lassa fever, 103, 114t
Latex allergies, 80–81
  prevention of in the workplace, 81
  products containing latex, 81t
  signs and symptoms of, 80
Laundry
  cleaning and disinfection of, 128
  Standard Precautions for, 101t, 104
Legionnaires' disease, 114t
Lensed instruments, cleaning and disinfection of, 127t
Leprosy, 114t
Leptospirosis, 114t
Lice, 114t
Lifestyle as risk for infection, 13
Linens, cleaning and disinfection of, 128
Lister, Joseph, 1
Listeria monocytogenes, 10t, 74t
Listeriosis, 114t
Luer lock, 44
Lyme disease, 114t
Lymphocytic choriomeningitis, 114t
Lymphogranuloma venereum, 114t

M

Mail-back programs, for disposal of medical sharps, 156
Malaria, 114t
Marburg virus, 103, 114t
Masks, 82, *83–84*
  Standard Precautions for, 101t
MBT *(Mycobacterium tuberculosis)*, 9, 10t
MDROs. *See* Multidrug resistant organisms (MDROs)
Meals for patients, 71, 76–78
  avoiding cross-contamination of food, 76, *76*
  cooking and serving, 77
  food storage, 76
  guidelines for preparation of, 76–77
  health department warnings, 78
  immunocompromised patients, 77–78

Meals for patients *(continued)*
microwaving, 77
overview of foodborne illness, 71, 72–75t
safety concerns if the power goes out, 77
shopping for, 71
Measles, 11t, 103, 117t
concerns for pregnant staff members regarding, 217t
congenital, 111t, 113t
defining and managing exposures to, 207
as indication for work restrictions, 213t
Measles vaccine, 200
Meatal care, 28
Medical devices. *See also specific devices*
categorization of, 122t
cleaning and disinfecting medical equipment before reuse, 26
cleaning and disinfecting respiratory equipment and supplies, 32
cleaning and disinfection of patient care equipment, 124–128, 127t
cleaning and disinfection of respiratory equipment and supplies, 31–32
cleaning enteral feeding equipment and supplies, 34
with engineered sharps injury protection feature, 131, 132–133t, 134
identification and usage of medical devices in the home, 121
as reservoir for for infectious agents, 9, 13–14, 170
Medical waste management, 131–162
cleaning blood spills in the home, 140
definitions, 134–135
labeling requirements, 139–140, *140*
medical waste, 134–135, 135
storage in the home care or hospice organization, 138–139
transport, 139
needle stick safety and prevention act
devices with engineered sharps injury protection feature, 131, 132–133t, 134
modification to the bloodborne pathogen standard, 131
problem-specific strategies for injury protection, 134t
recommendations of EPA and OSHA, 131
regulated waste, 135
regulations, 131
segregation of waste
general waste, 135
medical waste, 135–136
sharps, 136–138
separating waste from clean equipment and supplies, 138
storage during transport to home care or hospice organization, 138
Melioidosis, 115t
Meningitis, 115t
defining and managing exposures to, 207
as indication for empiric precautions, 106t
Meningococcal infection, 115t
Menningococcal infection, as indication for work restrictions, 213t
Methicillin-resistant *Staphylococcus aureus* (MRSA), 8, 9, 91
animals as carriers of, 69–71
Microorganisms
chain of infection, 8, 8f
common pathogenic agents, 10–11t
identifying the cause of outbreaks, 187–188
microscopic examination of, 9
multidrug resistance (*See* Multidrug resistant organisms (MDROs))
normal flora, 7–8
reservoirs of infection for, 9
resident, 15–16
transient, 16
Midline catheters, 39
infections related to, CDC guidelines for prevention of, 62
Minimal inhibitory concentration (MIC), 90
Molluscum contagiosusm, 115t
Monkeypox, 115t
MRSA. *See* Methicillin-resistant *Staphylococcus aureus* (MRSA)
Mucocutaneous herpes simplex, 113t
Multidrug-resistant organisms (MDROs), 89–96
animals as carriers of, 69–71
blood cultures, 90
CDC Guideline for Isolation Precautions, 89, 93–96, 95t
contamination of hme environment by, 96
earliest reports of, 89
education of staff, 96
epidemiology of, 91–92
gram negative, 90
gram positive, 90
identification of, 89–90
identifying patients with, 92–94
use of surveillance cultures, 92–93
Isolation Precautions, 115t, 118t
line listing data elements, 93–94, 93t
management of MDROs, 94–96, 95t
mechanisms of resistance, 91
penicillin, 89
risk of in ICUs, 91–92
susceptibility testing, 90, 91t
term defined, 89
transmission of, 90–91
in home and hospice, 92
types of antibiotics used to treat, 90
types of bacteria, 10t, 89–90

vancomycin-resistant *staphylococcus aureus (VRSA)*, 90–91, 91–92
in vitro resistance, 91
in vivo resistance, 91
wound cultures, 90
Mumps, 115t
as indication for work restrictions, 213t
Mumps vaccine, 200
Mycobacteria infection, nontuberculosis, 115t
Mycobacterium tuberculosis (MBT), 9, 10t
Mycoplasm pneumonia, 115t

N

National Association for Home Care (NAHC), 4
National Fire Protection Association (NFPA), classification of flammable liquids and foam, 21
National Institute for Occupational Safety and Health (NIOSH), protection from latex exposure and allergy, 81
National Nosocomial Infection Surveillance (NNIS) study, 2, 38, 163–164
definitions for hospital acquired infections, 166
definitions of nosocomial infections, 171
National Patient Safety Foundation (NPSF), grants related to home care safety, 221
National Phlebotomy Association, 126
Necrotizing enterocolitis, 115t
Necrotizing fasciitis, 118t
Needle destruction devices, 156
Needleless intravascular devices, 44, 131
Needle stick safety and prevention, 79. *See also* Sharps
devices with engineered sharps injury protection feature, 131, 132–133t, 134
modification to the bloodborne pathogen standard, 131
problem-specific strategies for injury protection, 134t
Standard Precautions for, 101t
Needlestick Safety and Prevention Act, 79, 131–134
Neisseria meningitidis disease, Droplet Precautions for, 103
Neonates. *See* Infants
NFPA (National Fire Protection Association), classification of flammable liquids and foam, 21
NIOSH (National Institute for Occupational Safety and Health), protection from latex exposure and allergy, 81
NNIS. *See* National Nosocomial Infection Surveillance (NNIS) study
Nocardiosis, 115t
Noncritical items, cleaning and disinfection of, 121, 122t
commonly used items, 125
laboratory supplies, 125–126
methods of sterilization and disinfection, 127t
provided to patients, 126
Nontunneled central venous catheters, 39–40
infections related to, CDC guidelines for prevention of, 63
Nonvascular access devices, 48–49
Normal flora, 7–8
Noroviruses, 112–113t, 115t
Norwalk agent gastroenteritis, 115t
Nosocomial infection
creating new standards for, 4
definitions of, 3, 4, 166, 171
earliest identification and analysis of, 1
epidemiology of, 2–3
extrinsic factors affecting rate of infection, 3
intrinsic factors affecting rate of infection, 3
NNIS definitions of, 166, 171
surveillance programs, 2, 38, 163–164
No-touch technique, wound care, 23, 24
NPSF (National Patient Safety Foundation), grants related to home care safety, 221
Nursing supply bag, 85–87
"bag technique," 86–87
contents of, 87
Nutrition. *See also* Infants, feeding
deficits in, and risk for infection, 13
enteral, 34
total parenteral, *47*, 48

O

Occupational health, 195–220
CDC recommendations for HIV postexposure prophylaxis, 219
defining and managing exposures, 205–212, *206*
to agents of bioterrorism, 208–209
to bloodborne pathogens, 209–212
to hepatitis A, 207
to measles, 207
to meningitis, 207
to pertussis, 207
to rubella, 207
to smallpox, 208–209
to tuberculosis, 207–208
to varicella, 206–207
designing a program for, 195–197, 196t
evaluation of services for, 197
health assessment and immunizations, 197
hepatitis B testing and vaccination, 195, 200–202, *201*
identifying health needs in planning, 195
influenza vaccine, 205
for pregnant staff members, 216, 217t
recordkeeping, 197
risk for infectious disease, 13, 195
tuberculosis assessment, 202–205
vaccine-preventable diseases, 197–202, *198*
diphtheria, 200
measles, 200

Occupational health *(continued)*
paying for testing and immunization, 198
pertussis, 200
polio, 200
preemployment health survey/immunization status, 198, *199*
rubella, 200
tetanus, 200
varicella, 198–200
work restrictions, indications for, 212–214t, 215–216
Occupational Safety and Health Administration (OSHA), 195
bloodborne pathogen regulations, 79, 123, 124, 131, 200–201, *201*
for HBV vaccine, 200–201, *201*
labeling requirements for medical waste, 139–140, *140*
log of occupational illnesses and injuries, 197
mandatory wearing of gloves, 80
selection of equipment to prevent needlestick injuries, 79
on TB skin testing in home care, 202
Ointments, antimicrobial or antiseptic, applied to catheter insertion site, 45
Orf, 115t
Ornithosis, 116t
Outbreak investigations, 187–193
case definition, development of, 189–191
communication to staff and patients, 193
community-acquired infections, 188
control of an outbreak, 191–192
data collection tool, *190*
definition of "outbreak," 187
documentation of, 192
epidemic curve—exit-site infections, 191, *191*
external assistance, 192–193
home care-acquired infections, 188
identifying the cause, 187–188
SARS epidemic, 187
steps in the investigation, 188–191, 189t
time frame of the outbreak, 191
why outbreaks should be investigated and reported, 187
Oxacillin, 38

P

Parainfluenza, 11t, 115t
Parasitic diseases, associated with zoonotic transmission, 70t
Parvovirus B19, 115t
concerns for pregnant staff members regarding, 216
Pasteur, Louis, 1
Pathogenic agents, 10–11t
Pathogenicity, 8, 8t
Patient care equipment, Standard Precautions for, 101t
Patient care practices, 2, 4, 15–35. *See also specific practices*
assessment of patient and home environment, 22
enteral therapy, 34
hand hygiene (*See* Hand hygiene)
post-mortem care, 34–35
prevention and control of influenza, 33–34
reusing equipment in the home, 26
urinary tract care (*See* Urinary tract care)
wound care (*See* Wound care)
Patient confidentiality, isolation precautions and, 106–107
Patient placement, Standard Precautions for, 101t
Patient resuscitation, Standard Precautions for, 101t
Pediatric patients, 65–68
cleaning and disinfecting clothing and linen, 66
cleaning and disinfecting diaper changing area, 66
cleaning and disinfecting "potty training" equipment, 66
cleaning of surfaces likely to be touched by, 65
diapering a child, *65*, 65–66
feeding
preparing infant formula
method, 67
storage of formula, 67–68
storage of human milk, 68
terminal sterilization, 67
thawing frozen milk, 68
preparing infants' bottles, 66–67
home infusion therapy, 43, 48
washing and disinfecting toys, 66
Pediculosis, 116t
as indication for work restrictions, 213t
Penicillin, 89
PEP (postexposure prophylaxis) for healthcare workers, 211–212
Periodic Payment System (PPS), 27
Peripherally inserted central catheter (PICC), 40. *See also* Home infusion therapy
catheter site care, 46
infections related to, CDC guidelines for, 62
infections related to, CDC guidelines for prevention of, 63
inserting catheter for, 43
for pediatric patients, 48
replacing, 43–44
selecting a catheter insertion site, 42–43
Peripheral venous catheters, 38–39, *39*, 48. *See also* Peripherally inserted central catheter (PICC)
infections related to, CDC guidelines for, 62
Personal protective equipment (PPE), 23, 79–86
bloodborne pathogen regulations, 79
criteria for type of PPE needed, 86t
donning, 82, *84*, 85
gloves
latex allergies and, 80–81
prevention of in the workplace, 81
products containing latex, 81t
microbial contamination of, 79

types of, 80
when to change or remove, 80
when to wear, 80
gowns, 82, *83–84*
infection control supplies, 87, *87*
masks, respiratory protection, eye protection, and face shields, 82, *83–84*
for needlestick prevention, 79
nursing supply bag, 85–87
"bag technique," 86–87
contents of, 87
"OSHA kit," 85
patient transport outside the home, 85
"PPE kit," 85
removing, 82, *84*, 85
resuscitation equipment, 85
staff member access to, 85
staff member barrier requirements, 86t
Standard Precautions for, 101t
training on location and use of, 85, 86t
use of, 79
Pertussis, 116t, 120t
defining and managing exposure to, 207
as indication for work restrictions, 213–214t
Pertussis vaccine, 200
PFAs (priority focus areas), 225
Pharmaceutical services, infection control in, 50–53
aseptic technique, 52
medication storage, 52
medication storage in patient's home, 52
multidose vials, 53
nursing care and administration of parenteral medications, 53
preparation of parenteral medication in the home, 52
storage and transport of parenteral medications, 52
USP-NF Chapter 797 requirements, 50, 51t, 57–59
Phenol derivatives, as antiseptic agents, 18t
Phenolic germicidal detergent solutions, 123
Phlebitis, infusion-related, 38–39, *39*
Pinworm infection, 116t
Plague, 116t
Plasmids, antibiotic resistance and, 91
Pneumococcal vaccine, 22
Pneumonia, 116t
definitions of for surveillance purposes, 172, 172t
in infants and young children, 118t
prevention, 30
indicators adapted for home and hospice care, *226*
performance indicators from CDC guidelines, *226*
Pocket infections, 37
Poliomyelitis, 116t
Polio vaccine, 200
Poliovirus, 11t
Portal of entry, 11
Portal of exit, 11
Postexposure prophylaxis (PEP) for healthcare workers, 211–212, 219–220
Post-mortem care, 34–35
Postpartum infection, 1
"Potty training" equipment, 66
Power failures during food preparation, 77
PPD (purified protein derivative) skin testing, 204–205
PPE. *See* Personal protective equipment (PPE)
Pregnancy, infections of concern to pregnant staff members, 216, 217t
Pressure ulcer, 111t
Prion disease, 116t
Priority focus areas (PFAs), 225
Protective Environment (PE), 100–101
Proteus sp., 10t, 27
Pseudomonas, 16, 90
as cause of BSIs, 38
as cause of urinary tract infections, 27
in ICU patients, 91
Pseudomonas aeruginosa, 9, 10t, 90
Pseudomonas cepacia, 90
Psittacosis, 116t
Puerperal fever, 1
Pulse oximeters, cleaning and disinfection of, 125
Purified protein derivative (PPD) skin testing, 204–205

Q

Q fever, 116t
Quality assurance practices, 58
Quality assurance program, 57
Quality control for pharmacies, 50, 51t, 57–59
Quaternary ammonium compounds, as antiseptic agents, 18t
Quinton catheter, 40

R

Rabies, 116t
Rash, as indication for empiric precautions, 106t
Rat-bite fever, 117t, 118t
Recapping a needle, 136
Recordkeeping. *See also* Data collection methods
occupational health, 197
Regulated medical waste, 135. *See also* Medical waste management
Relapsing fever, 117t
Rental of reusable equipment, 126
Reservoir of infection, 9
Resistance, 8t, 9
Resource guidelines for cleaning and disinfection, 121
Respiratory equipment and supplies, cleaning and disinfection of, 31–32
Respiratory Hygiene/Cough Etiquette, 100, 101t
for SARS patients, 102–103

Respiratory protection, 82, *83–84*
Respiratory syncytial virus, 11t, 117t
Respiratory therapy, 30–33
  breathing circuits, 30
  cleaning and disinfecting
    inner tracheal cannula, 31–32
    respiratory equipment and supplies, 32
    tracheal suction catheters, 31
  measures to prevent respiratory infection, 32–33
  mechanically assisted ventilation, 30
  patient education for prevention of respiratory tract infection, 33
  preventing aspiration, 33
  suctioning of respiratory tract secretions, 30–31
Respiratory tract infections, 117t
  as indication for empiric precautions, 106t
Responsibilities, inspection control practitioner, *224*
Resuscitation equipment, 85
Reye's syndrome, 117t
Rheumatic fever, 117t
Rhinovirus 39, 11t
Rickettsial fevers, 117t
Rickettsialpox, 117t
Ringworm, 117t, 118t
Risk factors for infection
  affecting rate of infection, 3
  associated with catheter-related infections during TPN, *47*
  extrinsic, 13–14
  intrinsic, 12–13
  reducing risk, 12–14
  risk factors by site, 173, 174t
  socioeconomic status and, 13
Ritter's disease, 117t
Rocky Mountain spotted fever, 117t
Rodents, transmission of infection by, 22
Rotavirus, 11t, 113t
Rubber tubing, cleaning and disinfection for, 127t
Rubella, 11t, 103, 117t
  concerns for pregnant staff members regarding, 217t
  congenital, 111t, 113t
  defining and managing exposure to, 207
  as indication for work restrictions, 214t
Rubella vaccine, 200
Rubeola, 103, 117t

S

SafeNeedleDisposal.org, 150
Salmonella, 8, 16, 74t, 113t
  as indication for work restrictions, 216
Salmonella choleraesuis, 10t
Salmonella paratyphi, 10t
Salmonella schottmuelleri, 10t
Salmonella typhi, 119t
Salmonellosis, 117t
SARS. *See* Severe Acute Respiratory Syndrome (SARS)
Scabies, 117t
  as indication for work restrictions, 214t
Scalded skin syndrome, 117t, 118t
Scales, cleaning and disinfection of, 125
Scarlet fever, 118t
Schistosomiasis, 117t
Semicritical items, cleaning and disinfection of, 121, 122t
Semmelweiss, Ignaz Philip, 1
Serratia sp., 10t, 27
Severe Acute Respiratory Syndrome (SARS), 117t, 187
  Isolation Precautions for, 100, 102–103, 117t
Sexually transmitted disease, 7, 11
Sharps, 136–138. *See also* Needle stick safety and prevention
  decontaminating before transport and disposal, 138
  disposal of, 138, 145, 146, 147
    helping your patients, 151
    at home, a book for young people with diabetes, 156–162
    questions for patients, 150
  recapping a needle, 136
  sharps container, 136–137, 158, 159, 160
  sharps-related injuries and factors related to, 136
  Standard Precautions for, 101t
  storage in the patient's home, 137
  storage in the staff member's possession, 137–138
  traveling with needles, 144
Sharps with engineered sharps injury protection (SEISP), 131
SHEA. *See* Society for Healthcare Epidemiologists of America (SHEA)
Shigatoxin-producing *E. coli* (STEC), 72t
Shigella, 8, 10t, 16, 74t, 113t
  as indication for work restrictions, 216
Shigellosis, 117t
Skin
  effects of antimicrobial soaps on, 19
  preparation of for home infusion therapy, 45
Skin infections
  as indication for empiric precautions, 106t
  as indication for work restrictions, 212, 215
Smallpox, 117t
  defining and managing exposure to, 208–209
  Isolation Precautions for, 102–103
  respiratory protection for, 82
Snow, John, 163
Soaps
  bar soap *vs.* liquid soap, 18
  plain *vs.* antimicrobial, for handwashing, 17–18, 18t

Society for Healthcare Epidemiologists of America (SHEA), 3–4
on the prevention of urinary tract infections, 27
recommendations for infection control programs, 222
Socioeconomic status, as risk factor for infection, 13
Sodium hypochlorite. *See* Bleach solution
Spaulding, E. H., 121
Specimen collection, urinary tract care and, 29
Spirillum minus disease, 118t
Spore-forming bacteria, 9
Standard Precautions, 100, 101t. *See also* Isolation precautions
Staphlococcal disease, 118t
skin infections, as indication for work restrictions, 212, 215
Staphylococcal food poisoning, 112t
Staphylococcal scalded skin syndrome, 117t
Staphylococcus aureus, 10t, 16, 74t, 90
as cause of BSIs, 38
methicillin-resistant, 8
Staphylococcus epidermidis, 8, 10t, 15, 90
as cause of IV catheter-related infections, 38, *39*
Staphylococcus pneumoniae, 10t
Sterilants, 123
Sterile technique
and clean technique, comparison of, 23
defined, 23
pharmacy sterile compounding requirements (USP and ASHP), 50, 51t, 57–59
procedures, 24
"Sterile to sterile" concept, 23
Sterilization, 123. *See also* Cleaning and disinfection
Stethoscopes, cleaning and disinfection of, 125
Streptobacillus monilliformis disease, 118t
Streptococcal disease, 118t
skin infections, as indication for work restrictions, 212, 215
Streptococcal infection, as indication for work restrictions, 214t
Streptococcus pneumoniae, 10t
Streptococcus pyogenes, 10t
Streptococcus salivarius, 10t
Streptococcus viridans, 7–8
Strongyloidiasis, 118t
Subclinical infection, 7
Suctioning of respiratory tract secretions, 30–31
Supplies, infection control, 87, *87*
Supply bags, cleaning and disinfection of, 125
Suprapubic catheter care, 28–29
Surfaces, cleaning and disinfection of, 127t
Surgical masks, 82, *83–84*
Surgical site infection surveillance, 170
Surveillance. *See* Infection surveillance
Susceptibility testing, 90, 91t
Susceptible host
extrinsic risk factors, 13–14
intrinsic risk factors, 12–13
Syphilis, 118t
Syringe Exchange Programs (SEP), 156

T

Taenia solium, 118t
Tapeworm infection, 118t
Terrorists, infectious agents likely to be used by, 208
Tetanus, 118t
Tetanus vaccine, 200
Textiles, Standard Precautions for, 101t
Thermometers, cleaning and disinfection of, 127t
Thrombophlebitis, 37
Tinea, 118t
"To Err is Human," 221, 224
Total parenteral nutrition (TPN), *47,* 48. *See also* Home infusion therapy
Towelettes, alcohol impregnated, 21
Toxic shock syndrome, 118t
Toxoplasmosis, 118t
concerns for pregnant staff members regarding, 217t
Tracheal cannula, cleaning and disinfecting, 31–32
Tracheal suction catheters
cleaning and disinfecting, 31
practices for reuse of, 30–31
Tracheostomy
frequency of tube changes, 32
stoma site care, 32
Trachoma, 118t
Training. *See* Education
Transmission of disease
airborne, 12
bloodborne, 7
carriers, 7
by contact, 11–12
in home care environment, 14
reducing the risk for, 12
by vector, 12
vehicle, 12
Transport
blood storage during, 49–50
Contact Precautions during patient transport, 105
decontamination of sharps before transport and disposal, 138
Droplet Precautions during patient transport, 104
of medical waste, 138, 139
patients infected with virulent organisms, 85
storage and transport of parenteral medications, 52
storage of equipment and supplies during, 129–130
use of personal protective equipment during patient

transport, 85
Trench mouth, 119t, 120t
Trichinosis, 119t
Trichomoniasis, 119t
Trichuriasis, 119t
Tricolsan, as antiseptic agent, 18t
Trivalent vaccine, 200
Trycophyton mentagrophytes, 11t
Tuberculoidal hospital disinfectants, 123–124
Tuberculosis, 119t
airborne precautions for, 82
concerns for pregnant staff members regarding, 217t
defining and managing exposure to, 207–208
as indication for work restrictions, 214t
Isolation Precautions for, 102
testing home-care staff for, 202–205
annual TB skin test, 204–205
PPD skin testing, 203–204
two-step testing, 204
Tularemia, 119t
Tunneled central venous catheters, 40, 63
Tunnel infections, 37
Typhoid fever, 119t
Typhus, 119t

U

United States Pharmacopoeia—National Formulary (USP-NF) Chapter 797 requirements, 50, 51t, 57–59
Universal Precautions, 100
Urinary tract care, 26–33
catheter-associated urinary tract infection (CAUTI), 26–27
catheter maintenance, 28
cleaning and disinfecting intermittent urethral catheters, 29
cleaning and disinfecting urine collection and tubing bags, 29
condom catheter drainage, 27
indwelling catheter insertion and replacement frequency, 27–28
indwelling catheter irrigation, 28
intermittent catheterization, 29
meatal care, 28
patient education for prevention of UTI, 29–30
suprapubic catheters, 28–29
Urinary tract infection (UTI), 119t
catheter-related, 26–27, 164
definitions of, 166, 166t
definitions of for surveillance purposes, 171, 172t
gender risk for, 13
patient education for prevention of, 29–30
prevention strategies, 27
risk for, 27
signs and symptoms of, 171
sources of infection, 27
symptoms of, 7
Urine collection tubing and bags, cleaning and disinfection of, 29
U.S. Postal Service, requirements for medical waste transport, 139
U.S. Public Health Service, on vaccines for children, 197
USP-NF Chapter 797 requirements, 50, 51t, 57–59
Utensils, eating, cleaning and disinfection of, 128
UTI. *See* Urinary tract infection (UTI)

V

Vaccination site care, 119t
Vaccine-preventable diseases, 197–202, *198*
Vaccines. *See* Immunizations
Vaccinia, 11t, 119t
Vancomycin-resistant enterococci (VRE), 8, 38, 91–92
animals as carriers of, 69
Vancomycin-resistant *staphylococcus aureus* (VRSA), 91–92
Varicella, 9, 110t, 120t
concerns for pregnant staff members regarding, 217t
defining and managing exposure to, 206–207
Varicella vaccine, 198–200
Varicella-zoster, 114t
Varicella zoster immune globulin (VZIG), 206
Variola. *See* Smallpox
Vector transmission of disease, 12
Vehicle transmission of disease, 12
Ventilation, mechanically assisted, risk of pneumonia and, 30
Viability, 8t, 9
Vibrio cholerae toxin, 74t
Vibrio parahaemolyticus, 74t, 113t
Vibrio vulnificus, 74t
Vincent's angina, 120t
Viral diseases
associated with zoonotic transmission, 70t
hemorrhagic fevers, respiratory protection, 82
respiratory, 120t
as indication for work restrictions, 214t
viral hemorrhagic fevers, 120t
Virulence, 8, 8t
Viruses, 9, 10–11t
VRE (vancomycin-resistant enterococci), 8
VZIG (varicella zoster immune globulin), 206

W

Warning labels for medical waste, 139–140, *140*
Waste. *See* Medical waste management
Waterless antiseptic agents, 17–18, 19
West Nile virus, 163

Whipworm disease, 119t
Whooping cough. *See* Pertussis
Work restrictions, indications for, 212–214t, 215–216
Wound care
  acute wounds, 22
  chronic wounds, 22
  clean technique *vs.* sterile technique, 23
  irrigating solutions
    maintenance of, 25, 25t
    preparation of, 25–26
  patient education relating to, 26
  procedures, 24–25
  selecting the "right" technique, 23–24, 24t
  surgical wounds, 22
  types of wounds cared for in the home, 22
Wound cultures, for bacteria, 90
Wound infections, 120t
  as indication for empiric precautions, 106t
  surveillance, 171

Y

Yersinia enterocolytica, 74t, 113t, 120t
Yersinia pestis, 116t
Yersinia pseudotuberculosis, 74t

Z

Zidovudine (ZVD), as postexposure prophylaxis for healthcare workers, 211
Zoonotic diseases, transmission of, 69–71, 70t
Zoster. *See* Herpes zoster
Zygomycosis, 120t